

TALENT Course 2018

Many-Body Perturbation Theory Calculations

胡柏山 (Baishan Hu)

北京大学物理学院 (Peking University, Beijing, China)

July 31, 2018 @ Xinxiang



Outline

I. Introduction

II. Theoretical framework and calculations

a) MBPT for closed-shell nuclei




b) MBPT (Q -box + folded diagrams) for open-shell nuclei

III. Summary and Outlook



Perturbation theory

takes us from a simple, exactly solvable (unperturbed) problem to a corresponding real (perturbed) problem



😊 The key formulas in perturbation theory are

$$\hat{H} = \hat{H}_0 + (\hat{H} - \hat{H}_0) = \hat{H}_0 + \hat{V} \quad \hat{H}_0 \Phi_n = E_n^{(0)} \Phi_n$$

😊 In MBPT the ambition is to include in H_0 as much “physics” as possible, so that V represents a “small” perturbation

History of MBPT

Key tool from 1950's to 1970's

- Rayleigh-Schrödinger perturbation theory
- G-matrix, Brueckner-Hartree-Fock method
- Valence-space shell-model interaction (Q-box + folded diagram)

Great depression in 1980's

- Depending on a starting energy parameter of G-matrix
- Poor convergence of the intermediate-state summations (tensor part)
- Intruder states

Today MBPT is coming back ...

- RSPT with soft potential ($V_{\text{low-}k}$, SRG, UCOM, OLS)
- Bogoliubov MBPT
- Auxiliary method (importance truncation, natural orbital basis)
- **Realistic Gamow shell model (Extended Kuo-Krenciglowa method)**

MBPT for closed-shell nuclei

$$\hat{H} = \sum_{i=1}^A \left(1 - \frac{1}{A}\right) \frac{\vec{p}_i^2}{2m} + \sum_{i<j}^A \left(\hat{V}_{NN,ij} - \frac{\vec{p}_i \cdot \vec{p}_j}{mA} \right) + \sum_{i<j<k}^A \hat{V}_{3N,ijk}$$

Rayleigh-Schrödinger perturbation theory

$$\hat{H} = \hat{H}_0 + (\hat{H} - \hat{H}_0) = \hat{H}_0 + \hat{V}$$

$$\hat{H}\psi_n = E_n\psi_n \quad \hat{H}_0\phi_n = E_n^{(0)}\phi_n$$

$$\Delta E = E_0 - E_0^{(0)}$$

$$\psi_0 = \sum_{m=0}^{\infty} [\hat{R}_0(E_0^{(0)})(\hat{V} - \Delta E)]^m \phi_0$$

$$\Delta E = \sum_{m=0}^{\infty} \langle \phi_0 | \hat{V} [\hat{R}_0(E_0^{(0)})(\hat{V} - \Delta E)]^m | \phi_0 \rangle$$

$$\text{where } \hat{R}_0 = \sum_{i \neq 0} \frac{|\phi_i\rangle\langle\phi_i|}{E_0^{(0)} - E_i^{(0)}}$$

$$E_0 = E_0^{(0)} + E_0^{(1)} + E_0^{(2)} + E_0^{(3)} + \dots$$

$$E_0^{(1)} = \langle \phi_0 | \hat{V} | \phi_0 \rangle$$

$$E_0^{(2)} = \langle \phi_0 | \hat{V} \hat{R}_0 \hat{V} | \phi_0 \rangle$$

$$E_0^{(3)} = \langle \phi_0 | \hat{V} \hat{R}_0 (\hat{V} - \langle \phi_0 | \hat{V} | \phi_0 \rangle) \hat{R}_0 \hat{V} | \phi_0 \rangle;$$



$$\psi_0 = \phi_0 + \psi_0^{(1)} + \psi_0^{(2)} + \dots$$

$$\psi_0^{(1)} = \hat{R}_0 \hat{V} | \phi_0 \rangle$$

$$\psi_0^{(2)} = \hat{R}_0 (\hat{V} - E_0^{(1)}) \hat{R}_0 \hat{V} | \phi_0 \rangle$$

Advantages in HF basis, compared with HO basis


- 😊 Faster convergence
- 😊 Some perturbation diagrams are cancelled out
- 😊 In HO basis, calculations could be $\hbar\omega$ dependent, while much less in HF basis



RSPT for closed-shell nuclei




- Perform a Hartree-Fock calculation
- HF state is chosen as a reference state H_0
- In the HF basis, make RSPT corrections
 - Energy up to 3rd order
 - Wave function up to 2nd order (One-body density)



$$E = E^{(0)} + E^{(1)} + E^{(2)} + E^{(3)} + \dots$$




$E^{(3)}$




$$\Psi = \Phi_{\text{HF}} + \Psi^{(1)} + \Psi^{(2)} + \dots$$

$\Psi^{(1)}$



$\Psi^{(2)}$



HF-RSPT calculations for ^{16}O with $\text{N}^3\text{LO-SRG}$, $N_{\text{shell}}=13$, $\hbar\omega=35$ MeV

Binding energy

	SRG flow parameter λ (fm^{-1})			
	1.5	2.0	2.5	3.0
Expt. [60]	-127.619	-127.619	-127.619	-127.619
SHF	-169.968	-133.169	-85.173	-44.102
PT2	-10.132	-29.497	-59.617	-88.326
PT3	-0.794	-1.931	-4.630	-7.339
SHF+PT2+PT3	-180.893	-164.597	-149.419	-139.767

3NF important !

B.S. Hu, F.R. Xu, *et al.*, PRC 94, 014303 (2017)

Point-proton rms radius

	SRG flow parameter λ (fm^{-1})			
	1.5	2.0	2.5	3.0
Expt.	2.581	2.581	2.581	2.581
SHF	2.098	2.096	2.201	2.345
PT2	0.011	0.011	-0.006	-0.042
$\Delta r_{\text{c.m.}}$	-0.067	-0.067	-0.070	-0.073
SHF+PT2+ $\Delta r_{\text{c.m.}}$	2.042	2.040	2.125	2.230

Inclusion of 3NF

$$\hat{H} = \sum_{i=1}^A \left(1 - \frac{1}{A}\right) \frac{\vec{p}_i^2}{2m} + \sum_{i<j}^A \left(\hat{V}_{NN,ij} - \frac{\vec{p}_i \cdot \vec{p}_j}{mA} \right) + \sum_{i<j<k}^A \hat{V}_{3N,ijk} = \sum_{i=1}^A \hat{H}_i^{(1)} + \sum_{i<j}^A \hat{H}_{ij}^{(2)} + \sum_{i<j<k}^A \hat{V}_{3N,ijk}$$

Normal ordering
with HF
reference state

$$\hat{H} = \sum_i \langle i | \hat{H}^{(1)} | i \rangle + \frac{1}{2} \sum_{ij} \langle ij | \hat{H}^{(2)} | ij \rangle + \frac{1}{6} \sum_{ijk} \langle ijk | \hat{V}_{3N} | ijk \rangle \quad \hat{H}_{\text{HF}}$$

$$+ \sum_{pq} \left(\langle p | \hat{H}^{(1)} | q \rangle + \sum_i \langle pi | \hat{H}^{(2)} | qi \rangle + \frac{1}{2} \sum_{ij} \langle pij | \hat{V}_{3N} | qij \rangle \right) : \hat{p}^\dagger \hat{q} :$$

$$+ \frac{1}{4} \sum_{pqrs} \left(\langle pq | \hat{H}^{(2)} | rs \rangle + \sum_i \langle pqi | \hat{V}_{3N} | rsi \rangle \right) : \hat{p}^\dagger \hat{q}^\dagger \hat{s} \hat{r} :$$

$$+ \frac{1}{36} \sum_{pqrstu} \langle pqr | \hat{V}_{3N} | stu \rangle : \hat{p}^\dagger \hat{q}^\dagger \hat{r}^\dagger \hat{u} \hat{s} :$$

➔ Discard residual 3B part:
NO2B approximation


$$\hat{H} = \hat{H}_{\text{HF}} + \hat{V}$$

$$\hat{V} = \frac{1}{4} \sum_{pqrs} \langle pq | \hat{W} | rs \rangle : \hat{p}^\dagger \hat{q}^\dagger \hat{s} \hat{r} : + \frac{1}{36} \sum_{pqrstu} \langle pqr | \hat{V}_{3N} | stu \rangle : \hat{p}^\dagger \hat{q}^\dagger \hat{r}^\dagger \hat{u} \hat{s} :$$


$$\hat{W} = \left(\langle pq | \hat{H}^{(2)} | rs \rangle + \sum_i \langle pqi | \hat{V}_{3N} | rsi \rangle \right) : \hat{p}^\dagger \hat{q}^\dagger \hat{s} \hat{r} :$$

ASG diagram expansion when 3NF is included







$E^{(3)}$: 56 terms (Derived for the first time)






ASG diagram expansion when 3NF is included











NO2B approximation:




$$\left(\langle pq | \hat{H}^{(2)} | rs \rangle + \sum_i \langle pqi | \hat{V}_{3N} | rsi \rangle \right)$$



$$\langle pqr | \hat{V}_{3N} | stu \rangle$$





NN -only N^3LO -SRG

Bare JISP16

$$+ \hat{V}_{3N}^{\text{ct}} = C_{3N} \delta^{(3)}(\vec{x}_1 - \vec{x}_2) \delta^{(3)}(\vec{x}_1 - \vec{x}_3)$$

$$N_{\text{shell}}=7, \hbar\omega=30 \text{ MeV}$$

HF-RSPT calculations with NO2B N^2LO_{sat} , $N_{\text{shell}}=13$, $\hbar\omega=22$ MeV

Binding energy


Nucleus	NN		$NN + 3N$		Expt.
	IM-SRG	HF-RSPT3	IM-SRG	HF-RSPT3	
${}^4\text{He}$	-27.36	-26.90	-29.09	-28.15	-28.30
${}^{14}\text{C}$	-133.38	-132.40	-104.16	-105.06	-105.29
${}^{22}\text{C}$	-179.72	-185.82	-114.79	-113.95	-119.18
${}^{16}\text{O}$	-171.95	-171.08	-124.13	-125.16	-127.62
${}^{22}\text{O}$	-242.37	-246.03	-160.02	-156.77	-162.03
${}^{24}\text{O}$	-265.58	-269.40	-166.26	-163.12	-168.97
${}^{40}\text{Ca}$	-610.89	-608.28	-311.47	-320.66	-342.05
${}^{48}\text{Ca}$	-783.40	-784.52	-376.75	-370.02	-416.00

B.S. Hu, T. Li, F.R. Xu, in preparation (2018)

Charge radius

Nucleus	NN		$NN + 3N$		Expt.
	IM-SRG	HF-RSPT1	IM-SRG	HF-RSPT1	
${}^4\text{He}$	1.64	1.66	1.69	1.75	1.6755(28)
${}^{14}\text{C}$	2.10	2.10	2.43	2.57	2.5025(87)
${}^{22}\text{C}$	2.05	2.02	2.53	2.63	—
${}^{16}\text{O}$	2.20	2.20	2.67	2.78	2.6991(52)
${}^{22}\text{O}$	2.13	2.10	2.66	2.75	—
${}^{24}\text{O}$	2.14	2.10	2.70	2.78	—
${}^{40}\text{Ca}$	2.61	2.58	3.40	3.49	3.4776(19)
${}^{48}\text{Ca}$	2.55	2.51	3.38	3.46	3.4771(20)

MBPT for open-shell nuclei



References

-  T.T.S. Kuo and E. Osnes, *Folded-Diagram Theory of the Effective Interaction in Atomic Nuclei*, Springer Lecture Notes in Physics, (Springer, Berlin, 1990) Vol. 364.
-  T.T.S. Kuo, et al. ,A Simple Method an Angular for Evaluating Momentum Goldstone Coupled Diagrams in Representation.ANNALS OF PHYSICS,132, 237-276
-  S.Y. Lee and K. Suzuki, *Phys. Lett. B* 91 (1980) 79; K. Suzuki and S.Y. Lee, *Prog. Theor. Phys.* 64 (1980) 2091.
-  M. Hjorth-Jensen, T.T.S. Kuo, and E. Osnes, *Realistic effective interactions for nuclear systems*, *Phys. Rep.*, 1995, 261, 125.

MBPT for open-shell nuclei

$$\begin{pmatrix} PHP & PHQ \\ \hline QHP & QHQ \end{pmatrix} \mathcal{H} = e^{-G} H e^G \begin{pmatrix} P\mathcal{H}P & P\mathcal{H}Q \\ \hline 0 & Q\mathcal{H}Q \end{pmatrix} \quad \begin{aligned} \hat{H} &= \hat{H}_0 + (\hat{H} - \hat{H}_0) \\ &= \hat{H}_0 + \hat{H}_1 \end{aligned}$$

\Rightarrow $Q\mathcal{H}P = 0$

$$H_{\text{eff}} = P\mathcal{H}P = P e^{-G} H e^G P$$

Kuo-Krenciglowa method (Folded-Diagram method)

$$V_{\text{eff}} = \hat{Q}(\varepsilon_0) - \hat{Q}'(\varepsilon_0) \int \hat{Q}(\varepsilon_0) + \hat{Q}'(\varepsilon_0) \int \hat{Q}(\varepsilon_0) \int \hat{Q}(\varepsilon_0) - \dots$$

$$V_{\text{eff}}^{(n)} = \hat{Q}(\varepsilon_0) + \sum_{k=1}^{\infty} \frac{1}{k!} \frac{d^k \hat{Q}(\varepsilon_0)}{d\varepsilon_0^k} \{V_{\text{eff}}^{(n-1)}\}^k$$

$$\hat{Q}(E) = P\hat{H}_1P + P\hat{H}_1Q \frac{1}{E - Q\hat{H}Q} Q\hat{H}_1P \quad \frac{1}{E - Q\hat{H}Q} = \sum_{n=0}^{\infty} \frac{1}{E - Q\hat{H}_0Q} \left(\frac{Q\hat{H}_1Q}{E - Q\hat{H}_0Q} \right)^n$$

☺ Factorization theorem: the core is separated out and only quantities relative to the core are concerned

☺ Q-box is made up of non-folded diagrams which are irreducible and valence linked

MBPT for open-shell nuclei

Bloch–Horowitz $H^{\text{BH}}(E_\lambda)P|\Psi\rangle = [PH_0P + P\hat{Q}(E_\lambda)P]P|\Psi\rangle = E_\lambda P|\Psi\rangle$

$$\hat{Q}(E) = P\hat{H}_1P + P\hat{H}_1Q \frac{1}{E - Q\hat{H}Q} Q\hat{H}_1P$$

BWPT

$$\frac{1}{E - Q\hat{H}Q} = \sum_{n=0}^{\infty} \frac{1}{E - Q\hat{H}_0Q} \left(\frac{Q\hat{H}_1Q}{E - Q\hat{H}_0Q} \right)^n$$

Kuo-Krenciglowa

degenerate model space

$$PH_0P = \varepsilon_0 P$$




$$V_{\text{eff}}^{(n)} = \hat{Q}(\varepsilon_0) + \sum_{k=1}^{\infty} \frac{1}{k!} \frac{d^k \hat{Q}(\varepsilon_0)}{d\varepsilon_0^k} \{V_{\text{eff}}^{(n-1)}\}^k$$

Extended Kuo-Krenciglowa

$$H_{\text{eff}}^{(n)} - E = PH_0P - E + \hat{Q}(E) + \sum_{k=1}^{\infty} \frac{1}{k!} \frac{d^k \hat{Q}(E)}{dE^k} \{H_{\text{eff}}^{(n-1)} - E\}^k$$

Advantages in HF basis, compared with HO basis

- 😊 **Faster convergence**
- 😊 **Some perturbation diagrams are cancelled out**
- 😊 **In HO basis, calculations could be $\hbar\omega$ dependent, while much less in HF basis**




K. Tsukiyama, *et al.*, PRC 85, 061304(R) (2012)






Diagrammatic expansion

S-box (1-body)

2nd




3rd







Q-box (2-body)

2nd



3rd



The pictures are taken from M. Hjorth-Jensen, *et al.*, Physics Reports 261 (1995) 125.

MBPT for open-shell nuclei

Realistic Gamow shell model Workflow

Bare forces:
Strong repulsion,
tensor force,
slow convergence

V_{low-k} ,
SRG,
OLS, ...

$$\langle ab|V_{osc}|cd\rangle \approx \sum_{\alpha \leq \beta}^N \sum_{\gamma \leq \delta}^N \langle ab|\alpha\beta\rangle \langle \alpha\beta|V_{low-k}|\gamma\delta\rangle \langle \gamma\delta|cd\rangle$$

$$\text{pp,nn: } \langle ab|\alpha\beta\rangle = \frac{\langle a|\alpha\rangle \langle b|\beta\rangle - (-1)^{J-j_a-j_b} \langle a|\beta\rangle \langle b|\alpha\rangle}{\sqrt{(1+\delta_{ab})(1+\delta_{\alpha\beta})}}$$

$$\text{pn: } \langle ab|\alpha\beta\rangle = \langle a|\alpha\rangle \langle b|\beta\rangle$$

HO/HF;
Gamow-Berggren
basis (a,b):
WS/GHF

$$\hat{H} = \hat{H}_0 + (\hat{H} - \hat{H}_0) = \hat{H}_0 + \hat{H}_1$$

$$\hat{Q}(E) = P\hat{H}_1P + P\hat{H}_1Q \frac{1}{E - Q\hat{H}Q} Q\hat{H}_1P$$

$$\frac{1}{E - Q\hat{H}Q} = \sum_{n=0}^{\infty} \frac{1}{E - Q\hat{H}_0Q} \left(\frac{Q\hat{H}_1Q}{E - Q\hat{H}_0Q} \right)^n$$



Up to third order

Valence-space effective
interactions
Complex shell model code
(Jacobi-Davidson, N. Michel)

Extended
Kuo-Krenciglowa
method (EKK)

$$H_{\text{eff}} = \begin{array}{c} j \\ | \\ \textcircled{\hat{Q}} \\ | \\ i \end{array} = \begin{array}{c} j \\ | \\ \textcircled{\hat{Q}} \\ | \\ i \end{array} - \begin{array}{c} j \\ | \\ \textcircled{\hat{Q}} \\ | \\ k \end{array} \int \begin{array}{c} j \\ | \\ \textcircled{\hat{Q}} \\ | \\ i \end{array} + \begin{array}{c} j \\ | \\ \textcircled{\hat{Q}} \\ | \\ l \end{array} \int \begin{array}{c} l \\ | \\ \textcircled{\hat{Q}} \\ | \\ k \end{array} \int \begin{array}{c} k \\ | \\ \textcircled{\hat{Q}} \\ | \\ i \end{array} - \dots$$

$$H_{\text{eff}} - E = PH_0P - E + \hat{Q}(E) + \sum_{k=1}^{\infty} \frac{1}{k!} \frac{d^k \hat{Q}(E)}{dE^k} \{H_{\text{eff}} - E\}^k$$



Gamow-Berggren basis

The wave function of a resonance with a peak at energy e_0 and a width γ

$$\Phi(e, \mathbf{r}) = \sqrt{\frac{\gamma/2}{\pi[(e - e_0)^2 + (\gamma/2)^2]}} \Psi(\mathbf{r})$$

Through the Fourier transform, we obtain the time evolution of the resonance

$$\Phi(t, \mathbf{r}) = \Psi(\mathbf{r}) e^{-i\tilde{e}_n t/\hbar} \quad \boxed{\tilde{e}_n = \frac{\hbar^2 k_n^2}{2m} = e_n - i\frac{\gamma_n}{2}} \quad t_{1/2} = \frac{\hbar \ln 2}{\gamma}$$

Gamow state: Complex energy G. Gamow, Z. Phys.51 (1928) 204


Berggren basis in complex- k plane, describing bound, resonance and scattering on equal footing.

T. Berggren, Nucl. Phys. A109 (1968) 265

Orthogonality and completeness

$$\delta(r - r') = \sum_n u_n(\tilde{e}_n, r) u_n(\tilde{e}_n, r') + \int_L d\tilde{e} u(\tilde{e}, r) u(\tilde{e}, r')$$

bound, resonance
scattering (discretized)




Gamow Hartree-Fock

Step 1: Solve the Hartree-Fock equations in HO representation using H_{int} ,

$$H_{\text{int}} = \sum_{i=1}^A \left(1 - \frac{1}{A}\right) \frac{\vec{p}_i^2}{2m} + \sum_{i<j}^A \left(V_{NN,ij} - \frac{\vec{p}_i \cdot \vec{p}_j}{mA} \right) + \sum_{i<j<k}^A V_{NNN,ijk}$$

Step 2: Extract the non-local HF potential $v(r,r')$

$$h_{ij}^{\text{HF}} = \langle i|t|j\rangle + \langle i|v|j\rangle = \langle i|t|j\rangle + \sum_{k=1}^A \langle ik|V|jk\rangle$$



Step 3: Obtain the radial wave function $u(r)/r$ in complex- k plane

$$u''(r) = \left[\frac{l(l+1)}{r^2} + v^{(\text{loc})}(r) - k^2 \right] u(r) + \int_0^{+\infty} v^{(\text{non-loc})}(r, r') u(r') dr'$$

$$u(\tilde{e}, r) \sim C_0 r^{l+1}, \quad r \rightarrow 0, \quad \tilde{e}_n = \frac{\hbar^2 k_n^2}{2m} = e_n - i \frac{\gamma_n}{2}$$

$$u(\tilde{e}, r) \sim C^+ H_{l\eta}^+(kr) + C^- H_{l\eta}^-(kr), \quad r \rightarrow +\infty.$$



Exterior complex scaling

$$\begin{aligned} \int_0^{+\infty} u(\tilde{e}, r)^2 dr &= \int_0^R u(\tilde{e}, r)^2 dr + (C^+)^2 \int_R^{+\infty} H_{l\eta}^+(kr)^2 dr \\ &= \int_0^R u(\tilde{e}, r)^2 dr + (C^+)^2 \int_0^{+\infty} H_{l\eta}^+(kR + kxe^{i\theta})^2 e^{i\theta} dx \end{aligned}$$

Outgoing solution at large distance

$$u_n(\tilde{e}_n, r) \sim O_l(k_n r) \sim e^{ik_n r}$$

Results of GHF



sp energies	^{16}O		^{22}O		^{24}O		^{28}O		MeV
	Re(E)	Im(E)	Re(E)	Im(E)	Re(E)	Im(E)	Re(E)	Im(E)	
$0s_{1/2}$	-48.858	0.000	-57.720	0.000	-59.313	0.000	-55.076	0.000	
$0p_{3/2}$	-22.735	0.000	-27.729	0.000	-28.132	0.000	-28.101	0.000	
$0p_{1/2}$	-13.863	0.000	-23.501	0.000	-22.669	0.000	-21.674	0.000	
$0d_{5/2}$	—	—	-3.251	0.000	-3.993	0.000	-6.687	0.000	
$1s_{1/2}$	—	—	-0.964	0.000	-2.374	0.000	-3.978	0.000	
$0d_{3/2}$	—	—	3.014	-0.626	2.312	-0.368	1.088	-0.081	

sp resonance

Order-by-order convergence in real-energy space



Bare NNLO_{opt}

$\hbar\omega=20$ MeV


$N_{\text{shell}}=12$

²²O core, *sd*-shell

SPEs Orbit	1st	2nd	3rd	Expt.
$\pi 0d_{5/2}$	-3.18	-12.01	-11.74	-13.26
$\pi 1s_{1/2}$	0.70	-7.95	-7.82	-10.88
$\pi 0d_{3/2}$	2.82	-7.15	-7.15	-
$\nu 1s_{1/2}$	-0.07	-4.42	-3.19	-2.73
$\nu 0d_{3/2}$	4.18	-0.12	1.31	1.26-0.2i




Oxygen isotopes from RGSM



Kondo, et al, (2016): ~ 0.110 MeV
Lunderberg, et al, (2012): ~ 0.005 MeV
Kohley et al, (2013): $4.5^{+11}_{-1.5}(\text{stat}) \pm 3$ (sys) ps

Fluorine isotopes from RGSM



Long-lived 4_1^+ isomer, $T_{1/2}=2.2(1)$ ms, Lepailleur, *et al.*, PRL110, 082502 (2013)

Gamow IM-SRG

**Hermite
(HO basis / real-energy HF)**

$$\langle a|H|b\rangle = \langle b|H|a\rangle^*$$

$$H(s) = U(s)H(0)U^\dagger(s)$$

$$U(s) \cdot U^\dagger(s) = U(s) \cdot U^{-1}(s) = 1$$

$$\eta(s) = \frac{dU(s)}{ds}U^\dagger(s) = -\eta^\dagger(s)$$



$$\frac{dH(s)}{ds} = [\eta(s), H(s)]$$

**Complex symmetric
(Gamow-Berggren basis)**


$$\langle \tilde{\alpha}|H|\tilde{\beta}\rangle = \langle \tilde{\beta}|H|\tilde{\alpha}\rangle^* = \langle \beta|H|\tilde{\alpha}\rangle$$

$$H(s) = U(s)H(0)U^T(s)$$


$$U(s) \cdot U^\dagger(s) = U(s) \cdot U^{-1}(s) = 1$$

$$\eta(s) = \frac{dU(s)}{ds}U^T(s) = -\eta^T(s)$$

Gamow IM-SRG vs RGSM



Gamow IM-SRG



Summary

- Develop HF-RSPT including the wave-function and three-body force corrections.
- Develop the third-order RGSM within HF basis
- Describe oxygen and fluorine isotopes

Outlook

- Derive the cross-shell effective interactions (*sdpf*-shell, ...)
- Reconciling the microscopic valence-space effective interactions with the nuclear reaction theory (GSM-RGM)

Collaborators:

Furong Xu (Peking University)

Qiang Wu (Peking University)

Tong Li (Michigan State University)

Zhonghao Sun (Oak Ridge National Laboratory)

Jianguo Li (Peking University)

Nicolas Michel (Michigan State University)

Gratitude:

Junchen Pei (Peking University)

Simin Wang (Michigan State University)

Thomas Papenbrock (Oak Ridge National Laboratory)

Gustav R. Jansen (Oak Ridge National Laboratory)

Luigi Coraggio (INFN-Naples)

James P. Vary (Iowa State University)

Ruprecht Machleidt (University of Idaho)

Marek Ploszajczak (GANIL)

Our group: Yuanzhuo Ma, Bo Dai, Sijie Dai, Yifang Geng, ...

Thank you for your attention!



Back up

A. Gamow Hartree-Fock

The HF single-particle state $|\alpha\rangle_{\text{HF}}$ can be expanded on HO basis $|p\rangle$,

$$|\alpha\rangle_{\text{HF}} = \sum_p D_{p\alpha} |p\rangle. \quad (2)$$

We diagonalize the Hartree-Fock one-body Hamiltonian in HO representation,

$$\langle p|h^{\text{HF}}|q\rangle = \langle p|t|q\rangle + \langle p|U|q\rangle = \langle p|t|q\rangle + \sum_{i=1}^A \sum_{rs} \langle pr|V|qs\rangle D_{ri}^* D_{si}. \quad (3)$$

After iterative solution of the HF equations, we can obtain the HF potential

$$\langle p|U|q\rangle = \sum_{i=1}^A \sum_{rs} \langle pr|V|qs\rangle D_{ri}^* D_{si}. \quad (4)$$

The the channel

$$\langle k|h^{\text{HF}}|k'\rangle = \left(1 - \frac{1}{A}\right) \frac{\hbar^2}{2m} k^2 \delta_{kk'} + \sum_{pq} \langle p|U_{\text{HF}}|q\rangle \langle k|p\rangle \langle q|k'\rangle \quad (5)$$

where $\langle q|k'\rangle$ is the HO wavefunction in complex-momentum space $\langle k|$. It's noting that

$$\hat{\rho}(\vec{r}) = \sum_{k=1}^A \delta^3(\vec{r} - \vec{r}_k) = \sum_{k=1}^A \frac{\delta(r - r_k)}{r^2} \sum_{lm} Y_{lm}^*(\hat{r}_k) Y_{lm}(\hat{r})$$

$$\hat{\rho}(\vec{r}) = \sum_K \sum_{n_1 l_1 j_1} \sum_{n_2 l_2 j_2} \sum_{m_j} R_{n_1 l_1}(r) R_{n_2 l_2}(r) \frac{-Y_{K0}^*(\hat{r})}{\sqrt{2K+1}}$$

$$\times \left\langle l_1 \frac{1}{2} j_1 \left| Y_K \right| l_2 \frac{1}{2} j_2 \right\rangle \langle j_1 m_j j_2 - m_j | K 0 \rangle$$

$$\times (-1)^{j_2 + m_j} a_{n_1 l_1 j_1 m_j}^\dagger a_{n_2 l_2 j_2 m_j}$$

$$\Psi = \Phi_0 + \Psi^{(1)} + \Psi_b^{(2)} + \Psi_c^{(2)},$$

$$\rho(\vec{r}) = \langle \Psi | \hat{\rho}(\vec{r}) | \Psi \rangle$$

$$= \langle \Phi_0 | \hat{\rho}(\vec{r}) | \Phi_0 \rangle + \langle \Phi_0 | \hat{\rho}(\vec{r}) | \Phi_0 \rangle \langle \Psi^{(1)} | \Psi^{(1)} \rangle + 2 \langle \Phi_0 | \hat{\rho}_N | \Psi_b^{(2)} \rangle + 2 \langle \Phi_0 | \hat{\rho}_N | \Psi_c^{(2)} \rangle + \langle \Psi^{(1)} | \hat{\rho}_N | \Psi^{(1)} \rangle$$

$$= \langle \Phi_0 | \hat{\rho}(\vec{r}) | \Phi_0 \rangle + \langle \Phi_0 | \hat{\rho}(\vec{r}) | \Phi_0 \rangle \langle \Psi^{(1)} | \Psi^{(1)} \rangle + 2\rho_a + 2\rho_b + \rho_{c_1} + \rho_{c_2},$$

$$\rho_a = \frac{1}{2} \sum_{h_1, h_2} \sum_{p_1, p_2, p_3} \frac{(-1)^{j_{h_1} + j_{h_2}} \sqrt{2j_{h_2} + 1}}{(\varepsilon_{h_1} - \varepsilon_{p_1})(\varepsilon_{h_1} + \varepsilon_{h_2} - \varepsilon_{p_2} - \varepsilon_{p_3})} \sum_J (-1)^J (2J + 1) \begin{Bmatrix} j_{h_1} & j_{p_1} & 0 \\ j_{h_2} & j_{h_2} & J \end{Bmatrix}$$

$$\times \langle (h_1 h_2) J | \hat{H} | (p_2 p_3) J \rangle \langle (p_2 p_3) J | \hat{H} | (p_1 h_2) J \rangle \langle h_1 || \rho || p_1 \rangle,$$



$$\begin{aligned}
& \langle ((n_1 l_1 j_1, n_2 l_2 j_2) J_1 T_1, n_3 l_3 j_3) J_2 T_2 | \hat{V} | ((n_4 l_4 j_4, n_5 l_5 j_5) J_3 T_3, n_6 l_6 j_6) J_2 T_2 \rangle \\
&= C_{3N} \sum_{L_1, S_1, L_2, S_2, L_3, S_3} \left\{ \begin{matrix} l_1 & \frac{1}{2} & j_1 \\ l_2 & \frac{1}{2} & j_2 \\ L_1 & S_1 & J_1 \end{matrix} \right\} \left\{ \begin{matrix} l_4 & \frac{1}{2} & j_4 \\ l_5 & \frac{1}{2} & j_5 \\ L_3 & S_3 & J_3 \end{matrix} \right\} \left\{ \begin{matrix} L_1 & S_1 & J_1 \\ l_3 & \frac{1}{2} & j_3 \\ L_2 & S_2 & J_2 \end{matrix} \right\} \left\{ \begin{matrix} L_3 & S_3 & J_3 \\ l_6 & \frac{1}{2} & j_6 \\ L_2 & S_2 & J_2 \end{matrix} \right\} \\
& \left\{ \delta_{S_1 S_3} \delta_{T_1 T_3} [1 - (-1)^{S_1 + T_1}] + \hat{S}_1 \hat{S}_3 \hat{T}_1 \hat{T}_3 \left\{ \begin{matrix} \frac{1}{2} & \frac{1}{2} & S_3 \\ \frac{1}{2} & S_2 & S_1 \end{matrix} \right\} \left\{ \begin{matrix} \frac{1}{2} & \frac{1}{2} & T_3 \\ \frac{1}{2} & T_2 & T_1 \end{matrix} \right\} [(-1)^{S_1 + T_1} + (-1)^{S_3 + T_3} - (-1)^{S_1 + T_1 + S_3 + T_3} - 1] \right\} \\
& \frac{1}{16\pi^2} \hat{l}_1 \hat{l}_2 \hat{l}_3 \hat{l}_4 \hat{l}_5 \hat{l}_6 \hat{L}_2^{-2} \begin{pmatrix} l_1 & l_2 & L_1 \\ 0 & 0 & 0 \end{pmatrix}_{\text{CG}} \begin{pmatrix} L_1 & l_3 & L_2 \\ 0 & 0 & 0 \end{pmatrix}_{\text{CG}} \begin{pmatrix} l_4 & l_5 & L_3 \\ 0 & 0 & 0 \end{pmatrix}_{\text{CG}} \begin{pmatrix} L_3 & l_6 & L_2 \\ 0 & 0 & 0 \end{pmatrix}_{\text{CG}} \\
& \int_0^\infty x^2 R_{n_1 l_1}(x) R_{n_2 l_2}(x) R_{n_3 l_3}(x) R_{n_4 l_4}(x) R_{n_5 l_5}(x) R_{n_6 l_6}(x) dx
\end{aligned}$$